

Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 111 / 229

CONTENTS

MODE BALL ABUTMENT INSTRUCTIONS FOR USE	112
MODE LOCATOR ABUTMENT INSTRUCTIONS FOR USE	120
MODE MULTI BASE ABUTMENT INSTRUCTIONS FOR USE	128
MODE ANGLED MULTI UNIT-N ABUTMENT INSTRUCTIONS FOR USE	137
MODE MULTI UNIT ABUTMENT INSTRUCTIONS FOR USE	145
MODE MULTI UNIT-N ABUTMENT INSTRUCTIONS FOR USE	152
MODE CEMENT RETAINED ABUTMENT INSTRUCTIONS FOR USE	160
MODE TEMPORARY ABUTMENT INSTRUCTIONS FOR USE	168
MODE TI-BASE and LONG TI-BASE ABUTMENT INSTRUCTIONS FOR USE	176
MODE FLEX TI-BASE AND ANGLED TI-BASE ABUTMENT INSTRUCTIONS FOR USE	185
MODE ABUTMENT AND PROSTHETIC SCREWS INSTRUCTIONS FOR USE	194
MODE HEALING ABUTMENT INSTRUCTIONS FOR USE	203
MODE SINUS COVER SCREWS INSTRUCTIONS FOR USE	213
MODE PROVO-S ENGAGED AND NON-ENGAGED INSTRUCTIONS FOR USE	223

Hazırlayan/ Prepared By Gonca Bakırcı

Al

Onaylayan/ Approverd By Saniye Özgür

Sayfa 111 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 112 / 229

MODE BALL ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Ball abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



			Narrow Platform		Reg	gular Platfo	orm
		Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm
	H0,5 H1,0 H2,0 Ball Abutment H4,0	H0,5	02.06.050.03		02.06.050.35		
		H1,0	02.06.01.03		02.06.01.35		
		H2,0	02.06.02.03		02.06.02.35		
		H3,0	02.06.03.03		02.06.03.35		
		H4,0	02.06.04.03		02.06.04.35		5
		H6,0 02.06.06.03 02.06.06.5				02.06.06.3	5

2. Intended Use

Mode Medikal Ball abutments are used to support dentures in order to fulfill the chewing function and to eliminate tooth deficiencies. One-piece Ball abutments are screwed to Mode Medikal implants and form a basis for overdenture structures.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.



KULLANMA KILAVUZU/ USER MANUAL

Ürün Adı: Dental İmplant Sistemi Üstyapıları

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 113 / 229

4. Indications

They are screwed directly to the intraosseous implants to support the removable overdentures.

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

Before the delivery of the prosthesis, the Ball abutment placed in the patient's mouth should be torqued with a torque value of 35 Ncm. Torque values greater than 35 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 114 / 229

- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

8. Compatibility Information

Metal and plastic matrices used with ball abutments are compatible with Rhein 83.

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Revision No: 07 Revision Date: 22.07.2024

Release Date:05.05.2020

Document No: TD.01/2.4.1

Page 115 / 229

Method Conditions Drying 121° C **Humidity temperature** Local (autoclave) 30 min

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

11. Procedure

pre-vacuum

11.1. Laboratory Procedure

Note: Screw the abutment to the analog during all laboratory procedures to preserve the abutment's attachment site to the implant.

- 1. Combine the impression coping with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Determine the abutment gingival height according to the amount of gingiva in the relevant region in the plaster model obtained.
- 4. Select the abutment suitable for the diameter of the dental implant used.
- 5. Connect the selected abutment with suitable diameter and gingival height to the analog in the plaster model with the help of screws.
- 6. Simply tighten the abutment screw by hand.
- 7. Plan your prosthesis in accordance with the selected abutment type and fabricate it with the preferred fabrication method.

9.2. Clinical Procedure

- 1. Remove the healing abutment located in the relevant area of the patient's mouth with the help of a screwdriver.
- 2. Select the abutment at the appropriate gingival height, insert it into the related implant in the mouth and screw the abutment with 35Ncm torque.
- 3. If the prosthesis and metal housing are to be combined in a laboratory environment, take the

impression with the relevant impression copings and custom impression tray.

- 4. At the delivery of the prosthesis, select the appropriate retention plastic and place it in the metal housing.
- 5. Place the prosthesis in the mouth and check the occlusal contacts, make the necessary adjustments.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 116 / 229

Oral and Prosthetic Care: Good oral hygiene is extremely important for the success of the attachment. Ball attachments should be thoroughly cleaned daily to prevent plaque buildup, and the patient should use a soft, nylon-bristled toothbrush and non-abrasive toothpaste to clean the abutments. Plastic retension materials are subject to wear as part of normal use and may need to be replaced. Patients should be asked to continue routine follow-up visits for hygiene and attachment function evaluation. Follow-up visits are recommended at 6-month intervals. Patients should be examined for signs of inflammation around the implant abutments and for implant mobility.

Placing and Removing Overdentures: The patient should be given instructions on how to properly place the overdenture. The patient should ensure that the prosthesis is correctly placed on the abutments before applying pressure. Using both hands, the patient should press both sides of the overdenture prosthesis to firmly seat it.

The patient should not bite the overdenture prosthesis into place; otherwise the applied force will cause improper wear of the abutments, including the plastic retainers in the overdenture. To remove the overdenture, the patient should place their thumbs under the edges of the overdenture and remove the prosthesis by pushing from both sides.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.



Ürün Adı: Dental İmplant Sistemi Üstyapıları

Revision No: 07 Revision Date: 22.07.2024

Release Date:05.05.2020

Document No: TD.01/2.4.1

Page 117 / 229

KULLANMA KILAVUZU/ USER MANUAL

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 117 | 229



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 118 / 229

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
8	Do Not Use Second Time	*	Do not expose to direct sunlight
i	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STEROLIZE	Do Not Sterilize For The Second Time

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Sog



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 119 / 229

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
₩	Name and Address of the implanting healthcare institution/provider
•••	Name and Address of the manufacturer
†i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 120 / 229

MODE LOCATOR ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Locator abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



		Narrow Platform		Reg	ular Platfo	orm
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm
	H0,5	02.07.050.03 02.07.05		2.07.050.03 02.07.050.35		5
14005	H1,0	02.07.01.03		02.07.01.35		
MODE Locator	H2,0	02.07.02.03		02.07.02.35		5
Abutment	H3,0	02.07.03.03		02.07.03.35		5
Abdiment	H4,0	02.07	.04.03	(2.07.04.3	5
	H6,0	02.07	.06.03	C	2.07.06.3	5

2. Intended use

Mode Medikal Locator abutments are used to support dentures in order to fulfill the chewing function and to eliminate tooth deficiencies. One-piece Locator abutments are screwed to Mode Medikal implants and form a basis for overdenture structures.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are screwed directly to the intraosseous implants to support the removable overdentures.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
AK .	Sarrye Ozgar



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 121 / 229

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

Before the delivery of the prosthesis, the Locator abutment placed in the patient's mouth should be torqued with a torque value of 35 Ncm. Torque values greater than 35 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 121 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 122 / 229

KULLANMA KILAVUZU/ USER MANUAL

- Minor bleeding
- Implant loss

8. Compatibility Information

Metal and plastic matrices used with Locator abutments are compatible with Zest Locator and Kerator.

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature	121° C	Local
(autoclave)	30 min	
pre-vacuum		



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Sayfa 123 | 229

Page 123 / 229

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

11. Procedure

11.1. Laboratory Procedure

Note: Screw the abutment to the analog during all laboratory procedures to preserve the abutment's attachment site to the implant.

- 1. Combine the impression coping with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Determine the abutment gingival height according to the amount of gingiva in the relevant region in the plaster model obtained.
- 4. Select the abutment suitable for the diameter of the dental implant used.
- 5. Connect the selected abutment with suitable diameter and gingival height to the analog in the plaster model with the help of screws.
- 6. Simply tighten the abutment screw by hand.
- 7. Plan your prosthesis in accordance with the selected abutment type and fabricate it with the preferred fabrication method.

9.2. Clinical Procedure

- 1. Remove the healing abutment located in the relevant area of the patient's mouth with the help of a screwdriver.
- 2. Select the abutment at the appropriate gingival height, insert it into the related implant in the mouth and screw the abutment with 35Ncm torque.
- 3. If the prosthesis and metal housing are to be combined in a laboratory environment, take the

impression with the relevant impression copings and custom impression tray.

- 4. At the delivery of the prosthesis, select the appropriate retention plastic and place it in the metal housing.
- 5. Place the prosthesis in the mouth and check the occlusal contacts, make the necessary adjustments.

Oral and Prosthetic Care: Good oral hygiene is extremely important for the success of the attachment. Locator attachments should be thoroughly cleaned daily to prevent plaque buildup, and the patient should use a soft, nylon-bristled toothbrush and non-abrasive toothpaste to clean the abutments. Plastic retension materials are subject to wear as part of normal use and may need to be replaced. Patients should be asked to continue routine follow-up visits for hygiene and attachment function evaluation. Follow-up visits are



Medical Devices Regulation (EU) 2017/745 CE Technical File **Product Name: Dental Implant System Superstructures** Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 124 / 229

recommended at 6-month intervals. Patients should be examined for signs of inflammation around the implant abutments and for implant mobility.

Placing and Removing Overdentures: The patient should be given instructions on how to properly place the overdenture. The patient should ensure that the prosthesis is correctly placed on the abutments before applying pressure. Using both hands, the patient should press both sides of the overdenture prosthesis to firmly seat it.

The patient should not bite the overdenture prosthesis into place; otherwise the applied force will cause improper wear of the abutments, including the plastic retainers in the overdenture. To remove the overdenture, the patient should place their thumbs under the edges of the overdenture and remove the prosthesis by pushing from both sides.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 125 / 229

responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 126 / 229

\triangle	Attention		Production Date
②	Do Not Use Second Time	类	Do not expose to direct sunlight
Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Song



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 127 / 229

VŖŲ.	Name and Address of the implanting healthcare institution/provider
***	Name and Address of the manufacturer
† i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Sayfa 127 | 229



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 128 / 229

MODE MULTI BASE ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Multi Base abutments are composed of two components. It consists of an abutment body and an abutment cover in different designs. Multi Base abutments are offered in two different angles, 17° and 30°.

Multi Base abutments offers a wide range of prosthetic solutions such as ball, Locator, bar-retained overdenture prostheses; and screwed fixed restorations.

Multi Base abutments, covers and copings are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



		Narrow Platform		Normal Platform		Wide Platform		
	Platform/H (mm)	Ø3.3mm	Ø3.7mm				Ø5.3mm	
17° Multi Base	H2,5	02.14.01.03		C	2.14.01.3	5	02.14.01.45	
Abutment	H3,5	02.14.02.03		02.14.02.35		02.14.02.45		
30° Multi Base	H3,5	02.14	.03.03	C	2.14.03.3	5		-
Abutment	H4,0	02.14	.04.03	C	2.14.04.3	5		-
Multi Base Cover		36.00.00.02						
Multi Base Ball	H1		02.16.00.01					
Cover	H2		02.16.00.02					
	H1	02.17.00.01						
Multi Base Locator Cover	H2	02.17.00.02						
Multi Base Ti Base Engaged Coping		02.07.00.03						
Multi Base Ti Base Non Engaged Coping		02.07.00.04						

2. Intended use

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Sig



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 129 / 229

Mode Medikal abutments are used for support in eliminating tooth deficiencies in order to fulfill the chewing function. Two-piece Mode Medikal Multi Base abutments and covers are attached to the implants with the help of screws and form the basis of prosthetic structures on implants such as screw retained fixed restorations and implant supported overdentures.

Multi Base Ti Base engaged and non engaged copings are supporting parts that provide a compatible interface with the Multi Base Cover by cementing the screwed final restoration or Geçici restoration in the laboratory environment.

3. Target patient group and intended user

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are screwed directly to the intraosseous implants with the help of screws in order to support the implantsupported prostheses. They provide support for screw retained fixed restorations or overdenture prostheses in edentulous or partially edentulous dental arches. Ball or Locator abutment cover is used to connect the removable overdentures with the implant. Multi Base abutments are used to connect bar-retained overdenture prostheses with the implant.

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medical implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

The abutment screw should be torqued with a torque value of 25 Ncm. Torque values greater than 25 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening. The prosthetic screws used in the Multi Base Abutment series should be torqued with a torque value of 15 Ncm.

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07 Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 130 / 229

KULLANMA KILAVUZU/ USER MANUAL

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications and side effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

8. Compatibility Information

Metal housing and retention plastics used with Multi Base Locator Cover are compatible with Zest Locator and Kerator. Metal and plastic matrices used with Multi Base Ball Cover are compatible with Rhein 83.

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page	131,	/ 229
------	------	-------

Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medical recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature	121° C	Local
(autoclave)	30 min	
pre-vacuum		

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

11. Procedure

11.1. Clinical Procedure

Impression

- 1. Multi Base abutments are selected in accordance with implant location, angle and gingival height.
- 2. The abutment body is placed in the appropriate position to the implant. The abutment carrier is flexible and can be bent if necessary.
- 3. The abutment body is fixed to the implant with a torque ratchet. The abutment body should be torqued with a torque value of 25 Ncm.
- 4. The abutment carrier is removed by turning.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı : Dental İmplant Sistemi Üstyapıları

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 132 / 229

KULLANMA KILAVUZU/ USER MANUAL

- 5. The abutment cover is placed on the abutment body and torqued with a standard hex wrench with a torque value of 25 Ncm.
- 6. It is recommended to check the fit of the abutment by radiography.
- 7. Implant impression is taken with open tray, closed tray or digital impression techniques.
- 8. If Geçici restoration will not be used, protective healing abutments are placed.
- 9. If a Geçici restoration is to be used, the Geçici restoration is placed on the abutments and the prosthetic screw is tightened by hand.





11.2. Laboratory Procedure

- 1. Combine the impression coping with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Produce the prosthetic infrastructure according to the preferred method. Infrastructure can be produced as casting, CAD/CAM or metal sintering.
- 4. Fabricate the final prosthesis according to the preferred laboratory technique.

11.3. Clinical Procedure

Restoration delivery

Screw Retained Fixed Restoration

- 1. Remove the healing abutment located in the relevant area of the patient's mouth in order to be able to try-in the prosthesis produced by the technician.
- 2. If the patient has a Geçici prosthesis with an occlusal screw, loosen the screws of the Geçici prosthesis with the help of a screwdriver and remove the Geçici prosthesis from the mouth.
- 3. Loosen the prosthetic screw with a screwdriver to remove the prosthesis on the abutment in the plaster model.
- 4. Remove the restoration from the model.
- 5. Place the prosthesis on the relevant implants on the patient in the same position as the model and tighten the prosthetic screw with the help of a screwdriver and a torque ratchet. The prosthetic screw should be torqued with a torque value of 15 Ncm.

Overdenture Restoration with Locator or Ball Attachment

- 1. Select the Multi Base abutment at the appropriate gingival height, place it in the mouth to the relevant implant and screw the abutment with 25 Ncm torque.
- 2. Insert the Multi Base Ball Cover or Multi Base Locator Cover and tighten with 25 Ncm torque value.
- 3. If the prosthesis and metal housing are to be processed in a laboratory environment, take an impression with the relevant impression coping and custom impression tray.
- 4. At the delivery of the prosthesis, select the appropriate retention plastic and place it in the metal housing.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



KULLANMA KILAVUZU/ USER MANUAL

Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07 Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 133 / 229

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

12. Magnetic Resonance Imaging (MRI) Safety Information

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medical products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 134 / 229

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention		Production Date
②	Do Not Use Second Time	类	Do not expose to direct sunlight



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 135 / 229

Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRUZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
吸	Name and Address of the implanting healthcare institution/provider



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 136 / 229

***	Name and Address of the manufacturer
†i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Shap



Medical Devices Regulation (EU) 2017/745 CE Technical File Product Name : Dental Implant System Superstructures Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 137 / 229

MODE ANGLED MULTI UNIT-N ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Multi Unit-N abutments are designed as a monoblok. Multi Unit-N abutments are offered in two different angles, 17° and 30°.

Multi Unit-N abutments offers a wide range of prosthetic Solutions.

Multi Unit-N abutments and copings are manufactured using Ti6Al4V-ELI (ASTM F 136) material.

10	E)
	U.
110	
VIII I	1
100	





		Narrow Platform		Regular Platform		orm
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm
	H1,5	02.14.10.03		02.14.20.03		
17° Anglad Multi	H2,5	02.14.05.03		02.14.05.35		
17° Angled Multi Unit-N Abutment	H3,5	02.14.06.03		02.14.06.35		
	H2,5	02.14.30.03		(02.14.40.0	3
30° Angled Multi	H3,5	02.14.07.03		(02.14.07.3	5
Unit-N Abutment	H4,5	02.14.08.03		02.14.08.35		5
Angled Multi Unit- N Ti Base Non Engaged Coping		02.07.00.02				

Hazırlayan/ Prepared By Gonca Bakırcı

Al

Onaylayan/ Approverd By Saniye Özgür

Sorg



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 138 / 229

2. Intended use

Mode Medikal abutments are used to support the replacement of missing teeth for the purpose of restoring chewing function. The one-piece Mode Medikal Angled Multi Unit-N abutments attach to implants with screws, serving as the foundation for screw-retained fixed restorations on implant-supported prosthetic structures.

The Angled Multi Unit-N Ti Base non-engaged copings are supportive parts that, when cemented in a laboratory setting, create a compatible interface with the Angled Multi Unit-N for screw-retained final or temporary restorations.

3. Target patient group and intended user

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are directly attached to endosseous implants with screws to serve as support for implant-supported prostheses. They provide support for fixed restorations in fully edentulous or partially edentulous dental arches.

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medical implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDICAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

The abutment screw should be torqued with a torque value of 25 Ncm. Torque values greater than 25 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening. The prosthetic screws used in the Multi Unit-N Abutment series should be torqued with a torque value of 15 Ncm.

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Medical Devices Regulation (EU) 2017/745 CE Technical File Product Name : Dental Implant System Superstructures Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 139 / 229

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications and side effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

8. Compatibility Information

All abutments and prosthetic screws can be used with the same abutment driver. Mode's two-piece implant series includes various types of abutments. Abbreviations and color codes indicating the compatibility of the implant and abutment platforms are present on the packaging.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2

9. Cleaning

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Medical Devices Regulation (EU) 2017/745 CE Technical File **Product Name: Dental Implant System Superstructures** Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 140 / 229

KULLANMA KILAVUZU/ USER MANUAL

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medical recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121° C 30 min	Local
pre-vacuum		

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

11. Procedure

11.1. Clinical Procedure

Impression

- 10. Angled Multi Unit-N abutments are selected in accordance with implant location, angle and gingival
- 11. The abutment body is placed in the appropriate position to the implant. The abutment carrier is flexible and can be bent if necessary.
- 12. The abutment body is fixed to the implant with a torque ratchet. The abutment body should be torqued with a torque value of 25 Ncm.
- 13. The abutment carrier is removed by turning.
- 14. It is recommended to check the fit of the abutment by radiography.
- 15. Implant impression is taken with open tray, closed tray or digital impression techniques.
- 16. If temporary restoration will not be used, protective healing abutments are placed.
- 17. If temporary restoration is to be used, the temporary restoration is placed on the abutments and the prosthetic screw is tightened by hand.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
A	Song



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 141 / 229

11.2. Laboratory Procedure

- 5. Combine the impression coping with the implant analogue.
- 6. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 7. Produce the prosthetic infrastructure according to the preferred method. Infrastructure can be produced as casting, CAD/CAM or metal sintering.
- 8. Fabricate the final prosthesis according to the preferred laboratory technique.

11.3. Clinical Procedure

Restoration delivery

Screw Retained Fixed Restoration

- 6. Remove the healing abutment located in the relevant area of the patient's mouth in order to be able to try-in the prosthesis produced by the technician.
- 7. If the patient has a Geçici prosthesis with an occlusal screw, loosen the screws of the Geçici prosthesis with the help of a screwdriver and remove the Geçici prosthesis from the mouth.
- 8. Loosen the prosthetic screw with a screwdriver to remove the prosthesis on the abutment in the plaster model.
- 9. Remove the restoration from the model.
- 10. Place the prosthesis on the relevant implants on the patient in the same position as the model and tighten the prosthetic screw with the help of a screwdriver and a torque ratchet. The prosthetic screw should be torqued with a torque value of 15 Ncm.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Sayfa 141 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 142 / 229

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medical products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 142 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 143 / 229

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
②	Do Not Use Second Time	*	Do not expose to direct sunlight
Ţ <u>i</u>	Please refer to the user manual	(Sa)	Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür




Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 144 / 229



Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
₩	Name and Address of the implanting healthcare institution/provider
***	Name and Address of the manufacturer
†i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Sig



Medical Devices Regulation (EU) 2017/745 CE Technical File Product Name: Dental Implant System Superstructures Ürün Adı: Dental İmplant Sistemi Üstvanıları

Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 145 / 229

MODE MULTI UNIT ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Multi Unit abutments are designed as one piece.

Mode Medikal Multi Unit abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



		Narrow Platform		Regular Platform			Wide Platform	
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm	Ø5.3mm	Ø6.0mm
	H0,5	02.05.00.03		02.05.01.35		02.05.00.45		
Multi-	H1,0	02.05.01.03		(02.05.02.3	5	02.05	.01.45
Unit	H2,0	02.05.02.03		(02.05.03.3	5	02.05	.02.45
Abutment	H3,0	02.05.03.03		02.05.04.35		02.05.03.45		
	H4,0	02.05.04.03		(02.05.05.35		02.05	.04.45

2. Intended Use

Mode Medikal abutments are used for support in eliminating tooth deficiencies in order to fulfill chewing function. It is attached to the two-piece Mode Medikal implants with the help of screws and forms the basis for screw retained fixed implant restorations.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are attached directly to the intraosseous implants with the help of screws in order to support the implantsupported prostheses. They provide support for fixed prostheses in edentulous or partially edentulous dental arches.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Revision No: 07 Revision Date: 22.07.2024

Release Date:05.05.2020

Document No: TD.01/2.4.1

Page 146 / 229

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

The Multi Unit abutment should be torqued to 25 Ncm. Torque values greater than 25 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening. The prosthetic screws used in the Multi Unit abutment series should be torqued with a torque value of 15 Ncm.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 147 / 229

• Implant loss

8. Compatibility Information

All abutments and prosthetic screws can be used with the same screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121° C 30 min	Local
pre-vacuum	30 11111	

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL Revision Page 148

Release Date:05.05.2020 Revision No: 07 Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 148 / 229

11. Procedure

11.1. Clinical Procedure

Impression

- 1. The Multi Unit abutment is selected in accordance with the gingival height.
- 2. The abutment is attached and tightened to the implant with a torque ratchet. The abutment should be torqued with a torque value of 25 Ncm.
- 3. It is recommended to check the fit of the abutment by radiography.
- 4. Implant impression is taken with open tray, closed tray or digital impression techniques.
- 5. If Geçici restoration will not be used, protective healing abutments are placed.
- 6. If a Geçici restoration is to be used, the Geçici restoration is placed on the abutments and the prosthetic screw is tightened by hand.

11.2. Laboratory Procedure

- 1. Combine the impression transfer with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Produce the prosthetic infrastructure according to the preferred method. Infrastructure can be produced as casting, CAD/CAM or metal sintering.
- 4. Fabricate the final prosthesis according to the preferred laboratory technique.

11.3. Clinical Procedure

Restoration Delivery

- 11. Remove the healing abutment located in the relevant area of the patient's mouth in order to be able to try-in the prosthesis produced by the technician.
- 12. If the patient has a Geçici prosthesis with an occlusal screw, loosen the screws of the Geçici prosthesis with the help of a screwdriver and remove the Geçici prosthesis from the mouth.
- 13. Loosen the prosthetic screw with a screwdriver to remove the prosthesis on the abutment in the plaster model.
- 14. Remove the restoration from the model.
- 15. Place the prosthesis on the relevant implants on the patient in the same position as the model and tighten the prosthetic screw with the help of a screwdriver and a torque ratchet. The prosthetic screw should be torqued with a torque value of 15 Ncm.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Hazırlayan/ Prepared By
Gonca Bakırcı

Onaylayan/ Approverd By
Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 149 / 229

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Savfa 149 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 150 / 229

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
②	Do Not Use Second Time	*	Do not expose to direct sunlight
Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 151 / 229

	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description		
† ?	Patient name or patient ID		
[31]	Date of implantation		
VĒV,	Name and Address of the implanting healthcare institution/provider		
***	Name and Address of the manufacturer		
F i	Information website for patients		
MD	Device name		
Hazırlayan/ Prepared By		Onaylayan/ Approverd By	



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 152 / 229

LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format
051	AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

MODE MULTI UNIT-N ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Multi Unit-N abutments are designed as one piece.

Mode Medikal Multi Unit-N abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



		Narrow Platform		Reg	gular Platfo	orm
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm
N 41±:	H1,5	02.16	.15.03	(02.16.15.3	5
Multi	H2,5	02.16	.25.03	(02.16.25.3	5
Unit-N Abutment	H3,5	02.16	.35.03	(02.16.35.3	5
Abutillelit	H4,5	02.16	.45.03	(02.16.45.3	5

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Sorg



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 153 / 229

2. Intended Use

Mode Medikal abutments are used for support in eliminating tooth deficiencies in order to fulfill chewing function. It is attached to the two-piece Mode Medikal implants with the help of screws and forms the basis for screw retained fixed implant restorations.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are attached directly to the intraosseous implants with the help of screws in order to support the implantsupported prostheses. They provide support for fixed prostheses in edentulous or partially edentulous dental arches.

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDICAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

The Multi Unit-N abutment should be torqued to 25 Ncm. Torque values greater than 25 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening. The prosthetic screws used in the Multi Unit-N abutment series should be torqued with a torque value of 15 Ncm.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07

Document No: TD.01/2.4.1

Revision Date: 22.07.2024

Page 154 / 229

KULLANMA KILAVUZU/ USER MANUAL

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

8. Compatibility Information

All abutments and prosthetic screws can be used with the same screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 155 / 229

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121° C	Local
pre-vacuum	30 min	

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

11. Procedure

11.1. Clinical Procedure

Impression

- 7. The Multi Unit-N abutment is selected in accordance with the gingival height.
- 8. The abutment is attached and tightened to the implant with a torque ratchet. The abutment should be torqued with a torque value of 25 Ncm.
- 9. It is recommended to check the fit of the abutment by radiography.
- 10. Implant impression is taken with open tray, closed tray or digital impression techniques.
- 11. If temporary restoration will not be used, protective healing abutments are placed.
- 12. If temporary restoration is to be used, the temporary restoration is placed on the abutments and the prosthetic screw is tightened by hand.

11.2. Laboratory Procedure

- 1. Combine the impression transfer with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Produce the prosthetic infrastructure according to the preferred method. Infrastructure can be produced as casting, CAD/CAM or metal sintering.
- 4. Fabricate the final prosthesis according to the preferred laboratory technique.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
A	Song



KULLANMA KILAVUZU/ USER MANUAL

Ürün Adı: Dental İmplant Sistemi Üstyapıları

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 156 / 229

11.3. Clinical Procedure

Restoration Delivery

- 16. Remove the healing abutment located in the relevant area of the patient's mouth in order to be able to try-in the prosthesis produced by the technician.
- 17. If the patient has a temporary prosthesis with an occlusal screw, loosen the screws of the temporary prosthesis with the help of a screwdriver and remove the Geçici prosthesis from the mouth.
- 18. Loosen the prosthetic screw with a screwdriver to remove the prosthesis on the abutment in the plaster
- 19. Remove the restoration from the model.
- 20. Place the prosthesis on the relevant implants on the patient in the same position as the model and tighten the prosthetic screw with the help of a screwdriver and a torque ratchet. The prosthetic screw should be torqued with a torque value of 15 Ncm.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 156 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 157 / 229

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By

Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 158 / 229

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention		Production Date
②	Do Not Use Second Time	*	Do not expose to direct sunlight
i	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	Ť	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Sog



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 159 / 229

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
Ų, T	Name and Address of the implanting healthcare institution/provider
***	Name and Address of the manufacturer
†i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 160 / 229

MODE CEMENT-RETAINED ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.

			Narrow	/ Platform	Regular Platform		tform	Wide Platform	
		Platform/I	d Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mi	m Ø5.2mm	ø5.3mm	Ø6.0mm
		H1,0	02.0	1.01.03		02.01.01	.35	02.01	01.45
	Profile	H2,0	02.0	1.02.03		02.01.02	.35	02.01	.02.45
	Abutment	H3,0	02.0	1.03.03		02.01.03	.35	02.01	03.45
		H1,0	02.1	3.01.03		02.13.01	.35	02.13	01.45
	Esthetic	H2,0	02.1	3.02.03		02.13.02	.35	02.13	02.45
	Abutment	H3,0	02.1	3.03.03		02.13.03	.35	02.13	03.45
		H1,0	02.0	8.01.03		02.08.01	.35	02.08	01.45
ALL	15° Esthetic	H2,0	02.0	8.02.03		02.08.02	.35	02.08	02.45
	Abutment	H3,0	02.0	8.03.03		02.08.03	.35	02.08	03.45
		H4,0	02.08.04.03		02.08.04.35		02.08.04.45		
		H1,0	02.0	9.01.03		02.09.01	.35	02.09	01.45
	25° Esthetic	H2,0	02.0	9.02.03		02.09.02	.35	02.09	02.45
	Abutment	H3,0	02.0	9.03.03		02.09.03	.35	02.09	03.45
		H4,0	02.0	02.09.04.03		02.09.04.35		02.09	04.45
	Premill Abutment		13.0	1.00.03		13.01.00	.35	13.01	00.45
			Narrow	Platform	Re	egular Plat	form	Wide P	latform
		Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm	Ø5.3mm	Ø6.0mm
		Abutment Diameter	Ø3,5mm	Ø4,0mm	Ø4,	5mm	Ø5,0mm	Ø5,0mm	Ø6.0mm
		H0,5	02.03.01.03	02.03.06.03	02.03	.06.35	02.03.11.35	02.03.01.45	02.03.06.45
	Direct	H1,0	02.03.02.03	02.03.07.03	02.03	.07.35	02.03.12.35	02.03.02.45	02.03.07.45
137	Abutment	H2,0	02.03.03.03	02.03.08.03	02.03	.08.35	02.03.13.35	02.03.03.45	02.03.08.45
		H3,0	02.03.04.03	02.03.09.03	02.03	.09.35	02.03.14.35	02.03.04.45	02.03.09.45

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 161 / 229

 H4,0
 02.03.05.03
 02.03.10.03
 02.03.10.35
 02.03.15.35
 02.03.05.45
 02.03.10.45

2. Intended Use

Mode Medikal abutments are used to support fixed implant supported prostheses in order to fulfill the chewing function and to eliminate tooth deficiencies. It is attached to the two-piece Mode Medikal implants with the help of abutment screws and forms a basis for cement retained prosthetic structures such as crowns and bridges.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are attached directly to the intraosseous implants with the help of abutment screws in order to support the implant-supported cement retained fixed prostheses.

	Indications		
Abutment type	Retention	Crown	Bridge
Profile Abutment	Cemented	✓	✓
Direct Abutment	Cemented	✓	✓
Esthetic Abutment	Cemented	✓	✓
15°/25° Esthetic Abutment	Cemented	✓	✓
Premill Abutment	Cemented	✓	✓

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDICAL can provide technical information. Please contact MODE MEDICAL sales representative.

Hazırlayan/ Prepared By
Gonca Bakırcı

Onaylayan/ Approverd By
Saniye Özgür

Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 162 / 229

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

Before the delivery of the prosthesis, the abutment placed in the patient's mouth should be torqued with a torque value of 35 Ncm. Torque values greater than 35 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

8. Compatibility Information

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Shap



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 163 / 229

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 1. Clean by brushing inside and outside with a brush under running water.
- 2. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 3. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121° C 30 min	Local
pre-vacuum		

Caution: Use devices immediately after sterilization. Do not store sterilizedk devices.

11. Procedure

11.1. Laboratory Procedure

Traditional Workflow in Implant Supported Fixed Dentures

Note: Screw the abutment to the analog during all laboratory procedures to preserve the abutment's attachment to the implant.

- 1. Combine the impression coping with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Determine the abutment gingival height according to the amount of gingiva in the relevant region in the plaster model obtained.
- 4. Select the abutment suitable for the diameter of the dental implant used.
- 5. Connect the selected abutment with suitable diameter and gingival height to the analog in the plaster model with the help of screws.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 164 / 229

- 6. Simply tighten the abutment screw by hand.
- 7. Plan your prosthesis in accordance with the selected abutment type and fabricate it with the preferred fabrication method.

Digital Workflow in Implant Supported Fixed Dentures

- 1. Digital prosthetic workflow can be carried out by using STL data obtained with intraoral scanners or by obtaining STL data of impressions taken with traditional methods with a desktop scanner.
- 2. The use of software that includes Mode implants in their library for prosthesis design and production on the digital platform will ensure that the internal geometry of the abutment and restoration connection is matching (eg Exocad DentalCAD).

11.2. Clinical Procedure for Implant Supported Fixed Prostheses

- 1. For the try-in of the prosthesis produced by the technician, remove the healing abutment located in the relevant area of the patient's mouth with the help of a screwdriver.
- 2. Loosen the abutment screw with a screwdriver to remove from the model.
- 3. Remove the abutment from the model.
- 4. Place the abutment on the patient's implant in the same position as the model and tighten the screw with the help of a screwdriver.
- 5. Perform the prosthesis try-in.
- 6. After try-in, remove the prosthesis from the patient's mouth and loosen the abutment screw with the help of a screwdriver.
- 7. Place the abutment on the corresponding analog on the model and tighten the abutment screw with the help of the screwdriver.
- 8. Before cementation of the prosthesis, tighten the screw of the abutment placed in the relevant area of the patient's mouth with a torque value of 35 Ncm.
- 9. It is recommended to check the fit of the abutment by radiography.
- 10. Cover the screw hole with a suitable material and cement the prosthesis with the preferred luting cement.

Warning: After the abutment screw is torqued, the abutment screw hole should be closed with suitable materials. Correct closure of the screw hole makes it possible to access the screw in cases where the abutment needs to be separated from the implant.

Abutment Modification

When necessary, abutments can be modified according to the patient's anatomy by the technician in the dental laboratory or by the dentist in the clinic.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Hazırlayan/ Prepared By
Gonca Bakırcı

Onaylayan/ Approverd By
Saniye Özgür

Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Revision No: 07 Revision Date: 22.07.2024

Release Date:05.05.2020

Document No: TD.01/2.4.1

Page 165 / 229

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred,

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 166 / 229

please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

130 13223 1: 2021 Symbols and Wealings Frepared According to the Standard							
Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol				
C € 2696	Notified Body Number	NON	Non Steril				
\triangle	Attention	M	Production Date				
②	Do Not Use Second Time	类	Do not expose to direct sunlight				
Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged				
REF	Reference Number	*	Keep away from contact with water				

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 167 / 229

	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
\F_	Name and Address of the implanting healthcare institution/provider
•••	Name and Address of the manufacturer
1	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code

Hazırlayan/ Prepared By
Gonca Bakırcı

Onaylayan/ Approverd By
Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 168 / 229



Explanation of unique device identifier as AIDC Format

AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

MODE TEMPORARY ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Temporary abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



		Narrow Platform		Reg	Regular Platform		Wide Platform	
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm	Ø5.3mm	Ø6.0mm
Multi Temporary Abutment				3	33.00.00.04	4		
Immediate	H1,5	02.18	.03.15	02.18.35.15		-		
Temporary Abutment	H3,0	02.18.03.03		02.18.35.03			-	
Temporary	H1,5	02.20	.03.15	02.20.35.15		02.20	.45.15	
Abutment Non-Engage	H3,0			-			02.20	.45.30
_	H1,5	02.19	.03.15	(02.19.35.1	5	02.19	.45.15
Temporary Abutment Engage	Н3,0			-			02.19	.45.30

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Sog




Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 169 / 229



2. Intended Use

Mode Medikal Temporary abutments are used to support Temporary prostheses to be used as an auxiliary tool in the treatment of tooth deficiencies. It attaches to the two-piece Mode Medikal implants with the help of screws and forms a basis for Temporary prosthetic structures such as crowns and bridges.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

3. Indications

They are attached directly to the intra-osseous implants with the help of screws in order to support the Temporary prosthesis on the implant. Temporary abutments are used as abutments in Temporary fixed prostheses.

	Indications		
Abutment type	Crown	Bridge	
Multi Temporary Abutment		✓	
Immediate Temporary Abutment	✓	✓	
Multi Unit-N Temporary Abutment	✓	✓	
Provo-S Temporary Abutment	✓	✓	
Temporary Abutment Engage	✓		
Temporary Abutment Non-Engage		✓	

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 170 / 229

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

It is contraindicated in patients with parafunctional habits and in individuals who do not have sufficient implant number and diameter to meet the occlusal forces.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

Mode Medikal abutments should only be used with Mode brand implants on the appropriate platform.

Before the delivery of the prosthesis, Immediate Temporary Abutment, Temporary Abutment Engage, Temporary Abutment Non-Engage screw should be torqued with a torque value of 35 Ncm. Torque values greater than 35 Ncm may damage the prosthetic parts. The Multi Temporary abutment should be torqued with a torque value of 15 Ncm. Torque values less than the recommended value may cause abutment loosening.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications and side effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 171 / 229

Implant loss

8. Compatibility Information

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 4. Clean by brushing inside and outside with a brush under running water.
- 5. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 6. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121° C	Local
pre-vacuum	30 min	

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
A	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 172 / 229

11. Procedure

- 1. Select the appropriate Temporary abutment for the planned Temporary prosthesis and connect it to the corresponding implant.
- 2. Check occlusal clearance and prepare out of the mouth for modification if necessary.
- 3. If preparation has been done, clean as specified in the cleaning and sterilization sections before inserting into the mouth and tighten the screw by hand.
- 4. Cement the clinical or laboratory prepared Temporary restoration to the abutment with a suitable material.
- 5. Drill a hole in the restoration for screw access. Loosen the screw with screwdriver and remove the restoration.
- 6. Finish and polish the restoration. Make the necessary occlusal adjustments.
- 7. Place the completed restoration in the mouth and torque the abutment screw with a ratchet to the appropriate torque.
- 8. After the abutment screw is torqued, close the abutment screw hole with suitable materials.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 173 / 229

Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Savfa 173 | 229



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 174 / 229

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
②	Do Not Use Second Time	*	Do not expose to direct sunlight
Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	Ť	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STENDIZE	Do Not Sterilize For The Second Time

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Sog



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 175 / 229

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
vīv, →	Name and Address of the implanting healthcare institution/provider
•••	Name and Address of the manufacturer
†i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Medical Devices Regulation (EU) 2017/745 CE Technical File Product Name: Dental Implant System Superstructures Ürün Adı: Dental İmplant Sistemi Üstvapıları

Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 176 / 229

MODE TI-BASE and LONG TI-BASE ABUTMENT INSTRUCTIONS FOR USE

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Ti-base and Long Ti-Base abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.

		Narrow Platform		Regular Platform		Wide Platform		
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm	Ø5.3mm	Ø6.0mm
	H0,7	17.00	.01.03	1	7.00.01.35	5	17.00	01.45
Ti-Base Cerec Abutment	H2,5	17.00	.02.03	1	17.00.02.3	5	17.00	02.45
Ti-Base	H0,7	16.00	.00.10	1	6.00.00.1	L	16.00	00.12
Engaged (Digital) Abutment	H2,5	16.00	.00.13	1	L6.00.00.1	1	16.00	00.15
Ti-Base Non-	H0,7	16.00	.00.04	1	16.00.00.0	5	16.00	00.06
Engaged (Digital) Abutment	H2,5	16.00	.00.07	1	16.00.00.08	3	16.00	00.09
Long Ti-Base Engaged	H0,7	16.90	.08.10	1	16.90.08.1	ı	16.90	08.12
(Digital) Abutment	H2,5	16.90	.08.13	1	16.90.08.14	1	16.90	08.15
Long Ti-Base	H0,7	16.90	.09.04	1	16.90.09.0	5	16.90	09.06
Non-Engaged (Digital) Abutment	H2,5	16.90	.09.07	1	16.90.09.08	3	16.90	09.09

2. Intended Use

Hazırlayan/ Prepared By
Gonca Bakırcı
Onaylayan/ Approverd By
Saniye Özgür

Sayfa 176 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 177 / 229

KULLANMA KILAVUZU/ USER MANUAL

Mode Medikal Ti-base and Long Ti-Base abutments are used to support fixed implant supported prostheses in order to fulfill the chewing function and to eliminate tooth deficiencies. It is attached to the two-piece Mode Medikal implants with the help of abutment screws and forms a basis for screw retained prosthetic structures such as crowns and bridges.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are attached directly to the intraosseous implants with the help of abutment screws in order to support the implant-supported cement or screw retained fixed prostheses.

Mode Medikal Ti-base and Long Ti-Base abutments are titanium bases that can also be used in personal abutment construction.

		Indica	ations
Abutment type	Retention	Crown	Bridge
Ti-Base Cerec Abutment	Screw	✓	
Ti-Base Engaged (Digital) Abutment	Screw	✓	
Long Ti-Base Engaged (Digital) Abutment	Screw	✓	
Ti-Base Non-Engaged (Digital) Abutment	Screw		✓
Long Ti-Base Non-Engaged (Digital) Abutment	Screw		√

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
AK	Solog



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 178 / 229

Mode Medical abutments should only be used with Mode brand implants on the appropriate platform.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDICAL can provide technical information. Please contact MODE MEDICAL sales representative.

Before the delivery of the prosthesis, the abutment placed in the patient's mouth should be torqued with a torque value of 35 Ncm. Torque values greater than 35 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

8. Compatibility Information

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7

Hazırlayan/ Prepared By

Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 179 / 229

Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2
Wide Platform	WP	Green	Ø5.3- Ø6.0

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 7. Clean by brushing inside and outside with a brush under running water.
- 8. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 9. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature	121° C	Local
(autoclave)	30 min	
pre-vacuum		

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

11. Procedure

11.1. Laboratory Procedure

Note: Screw the abutment to the analog during all laboratory procedures to preserve the abutment's attachment to the implant.

- 1. Combine the impression coping with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
A	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 180 / 229

- 3. Determine the abutment gingival height according to the amount of gingiva in the relevant region in the plaster model obtained.
- 4. Select the abutment suitable for the diameter of the dental implant used.
- 5. Connect the selected abutment with suitable diameter and gingival height to the analog in the plaster model with the help of screws.
- 6. Simply tighten the abutment screw by hand.
- 7. Plan your prosthesis with screw or cement retained in accordance with the selected abutment type and produce it with the preferred production method.
- 8. Place the fabricated restoration to the analog before cementing it to the Ti-base abutment.
- 9. Seal the screw channel with wax.
- 10. Apply a self-curing resin cement to the abutment. Use the resin cement in accordance with the manufacturer's instructions.
- 11. Place the restoration on the Ti-base abutment.
- 12. Immediately clean any cement overflowing from the abutment. Polish the joint after the cement has hardened.

Digital Workflow in Implant Supported Fixed Dentures

- 1. Digital prosthetic workflow can be carried out by using STL data obtained with intraoral scanners or by obtaining STL data of impressions taken with traditional methods with a desktop scanner.
- 2. The use of software that includes Mode implants in their library for prosthesis design and production on the digital platform will ensure that the internal geometry of the abutment and restoration connection is matching (eg Exocad DentalCAD).

11.2. Clinical Procedure for Implant Supported Fixed Prostheses

- 11. For the try-in of the prosthesis produced by the technician, remove the healing abutment located in the relevant area of the patient's mouth with the help of a screwdriver.
- 12. Loosen the abutment screw with a screwdriver to remove from the model.
- 13. Remove the abutment from the model.
- 14. Place the abutment on the patient's implant in the same position as the model and tighten the screw with the help of a screwdriver.
- 15. Perform the prosthesis try-in.
- 16. After try-in, remove the prosthesis from the patient's mouth and loosen the abutment screw with the help of a screwdriver.
- 17. Place the abutment on the corresponding analog on the model and tighten the abutment screw with the help of the screwdriver.
- 18. Before cementation of the prosthesis, tighten the screw of the abutment placed in the relevant area of the patient's mouth with a torque value of 35 Ncm.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 181 / 229

- 19. It is recommended to check the fit of the abutment by radiography.
- 20. Cover the screw hole with a suitable material and cement the prosthesis with the preferred luting cement.

Warning: After the abutment screw is torqued, the abutment screw hole should be closed with suitable materials. Correct closure of the screw hole makes it possible to access the screw in cases where the abutment needs to be separated from the implant.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 182 / 229

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

	100 101 10 10 10 10 10 10 10 10 10 10 10				
	Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol	
-	C € ₂₆₉₆	Notified Body Number	NON	Non Steril	

Hazırlayan/ Prepared By Gonca Bakırcı

Saniye Özgür

Onaylayan/ Approverd By



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 183 / 229

\triangle	Attention		Production Date
2	Do Not Use Second Time	类	Do not expose to direct sunlight
Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	于	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Song



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 184 / 229

[31]	Date of implantation
吸	Name and Address of the implanting healthcare institution/provider
~	Name and Address of the manufacturer
1	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı

Al



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 185 / 229

MODE FLEX TI-BASE AND ANGLED TI-BASE ABUTMENT **INSTRUCTIONS FOR USE**

1. Product Description

The Mode Medikal prosthesis series consists of abutments in different types of diameters, lengths and platforms that are used in the restoration of Mode dental implants. They are available in a variety of shapes and sizes to suit the patient's needs. These instructions for use are valid for Mode Medikal's products specified in section 3.

Mode Medikal Ti-base abutments are manufactured using Ti6Al4V-ELI (ASTM F 136) material.



		Narrow	Platform	Regular Platform		orm
	Platform/H (mm)	Ø3.3mm	Ø3.7mm	Ø4.1mm	Ø4.7mm	Ø5.2mm
	H0,7	17.00	.00.10	-	17.00.00.1	1
Flex Ti-Base Engaged (Digital) Abutment	H2,5	17.00	.00.13	<u>-</u>	17.00.00.1	4
	H0,7	17.00	.00.04	-	17.00.00.0	5
Flex Ti-Base Non-Engaged (Digital) Abutment	H2,5	17.00	.00.07	<u>-</u>	17.00.00.0	8
20° Angled Ti- Base Engaged (Digital) Abutment	H0,7	03.15.00.03		(03.15.00.3	5
	H2,5	03.15	.01.03	(3.15.01.3	5
20° Angled Ti- Base Non-	H0,7	03.15	.02.03	(03.15.02.3	5

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024 Page 186 / 229



Engaged (Digital) Abutment H2,5 03.15.03.03 03.15.03.35	Engaged (Digital) Abutment	H2,5	03.15.03.03	03.15.03.35
---	----------------------------------	------	-------------	-------------

2. Intended Use

Mode Medikal Ti-base abutments are used to support fixed implant supported prostheses in order to fulfill the chewing function and to eliminate tooth deficiencies. It is attached to the two-piece Mode Medikal implants with the help of abutment screws and forms a basis for screw retained prosthetic structures such as crowns and bridges.

3. Target Patient Group And Intended User

Mode dental implants are intended to be used in patients with complete or partial edentulism who have completed their growth and development and do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to safely and properly use Mode dental implants.

4. Indications

They are attached directly to the intraosseous implants with the help of abutment screws in order to support the implant-supported cement or screw retained fixed prostheses.

Mode Medikal Ti-base abutments are titanium bases that can also be used in personal abutment construction. Flex Ti Base is used in the anterior regions for aesthetic reasons due to its cut surface. Angled Ti Base abutments, on the other hand, are used in implants that are loaded at an angle.

		Indica	ations
Abutment type	Retention	Crown	Bridge
Flex Ti-Base Engaged (Digital) Abutment	Screw	✓	
Angled Ti-Base Engaged (Digital) Abutment	Screw	✓	
Flex Ti-Base Non-Engaged (Digital)	Screw		✓
Abutment			

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 187 / 229

Angled Ti-Base Non-Engaged (Digital)	Screw	✓
Abutment		

5. Contraindications

Its use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys.

For implant treatment contraindications, refer to the relevant Mode Medikal implant instructions for use.

6. Warnings and Precautions

The patient may swallow or aspirate the prosthetic parts. Care must be taken when working inside the mouth.

Mode Medical abutments should only be used with Mode brand implants on the appropriate platform.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

Before the delivery of the prosthesis, the abutment placed in the patient's mouth should be torqued with a torque value of 35 Ncm. Torque values greater than 35 Ncm may damage the prosthetic parts. Torque values less than the recommended value may cause abutment loosening.

Abutments are disposable devices.

Mode Medikal products should be used according to the instructions for use provided by the manufacturer. It is the physician's responsibility to decide whether the device is suitable for use according to these instructions and for the individual patient situation.

For a successful implant treatment, it is important that the dentist and dental technician work in harmony.

In order for the implant treatment to be successful in the long term, it is important that the patient's routine controls and oral hygiene training are given after the treatment is completed.

7. Complications And Side Effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Swallowing or aspiration of prosthetic components
- Peri-implantitis
- Implant component or prosthesis failure/fracture
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 188 / 229

Implant loss

8. Compatibility Information

All abutments can be used with the same abutment screwdriver.

There are different shapes of abutments in the Mode two-piece implant series. Abbreviations and color codes indicating the compatibility of implant and abutment platforms are available on the packages.

Implant Platform	Abbreviation	Color Code	Implant Diameter
Narrow Platform	NP	Yellow	Ø 3.3 – Ø3.7
Regular Platform	RP	Blue	Ø4.1 – Ø4.7 – Ø 5.2

9. Cleaning

Mode Medikal abutments and parts are delivered unsterile. Before placing the products in the patient's mouth, they must be disassembled, cleaned and sterilized.

Mode Medikal recommends the following for cleaning and sterilization of abutments before use.

- 10. Clean by brushing inside and outside with a brush under running water.
- 11. The pre-treated product can be cleaned by hand, ultrasonic cleaner, or an automatic cleaning method.
- 12. When selecting the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

10. Sterilization

Mode Medikal abutments and parts are delivered unsterile. Mode Medikal recommends the following for sterilization of abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121° C 30 min	Local
pre-vacuum		

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



KULLANMA KILAVUZU/ USER MANUAL

Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 189 / 229

11. Procedure

11.1. Laboratory Procedure

Note: Screw the abutment to the analog during all laboratory procedures to preserve the abutment's attachment to the implant.

- 1. Combine the impression coping with the implant analogue.
- 2. Obtain the plaster working model by using soft gingival silicone around the analogues.
- 3. Determine the abutment gingival height according to the amount of gingiva in the relevant region in the plaster model obtained.
- 4. Select the abutment suitable for the diameter of the dental implant used.
- 5. Connect the selected abutment with suitable diameter and gingival height to the analog in the plaster model with the help of screws.
- 6. Simply tighten the abutment screw by hand.
- 7. Plan your prosthesis with screw or cement retained in accordance with the selected abutment type and produce it with the preferred production method.
- 8. Place the fabricated restoration to the analog before cementing it to the Flex Ti-base and Angled Ti-base abutment.
- 9. Seal the screw channel with wax.
- 10. Apply a self-curing resin cement to the abutment. Use the resin cement in accordance with the manufacturer's instructions.
- 11. Place the restoration on the Flex Ti-base and Angled Ti-base abutment.
- 12. Immediately clean any cement overflowing from the abutment. Polish the joint after the cement has hardened.

Digital Workflow in Implant Supported Fixed Dentures

- 3. Digital prosthetic workflow can be carried out by using STL data obtained with intraoral scanners or by obtaining STL data of impressions taken with traditional methods with a desktop scanner.
- 4. The use of software that includes Mode implants in their library for prosthesis design and production on the digital platform will ensure that the internal geometry of the abutment and restoration connection is matching (eg Exocad DentalCAD).

11.2. Clinical Procedure for Implant Supported Fixed Prostheses

- 21. For the try-in of the prosthesis produced by the technician, remove the healing abutment located in the relevant area of the patient's mouth with the help of a screwdriver.
- 22. Loosen the abutment screw with a screwdriver to remove from the model.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Medical Devices Regulation (EU) 2017/745 CE Technical File Product Name: Dental Implant System Superstructures Ürün Adı: Dental İmplant Sistemi Üstvanıları

Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 190 / 229

- 23. Remove the abutment from the model.
- 24. Place the abutment on the patient's implant in the same position as the model and tighten the screw with the help of a screwdriver.
- 25. Perform the prosthesis try-in.
- 26. After try-in, remove the prosthesis from the patient's mouth and loosen the abutment screw with the help of a screwdriver.
- 27. Place the abutment on the corresponding analog on the model and tighten the abutment screw with the help of the screwdriver.
- 28. Before cementation of the prosthesis, tighten the screw of the abutment placed in the relevant area of the patient's mouth with a torque value of 35 Ncm.
- 29. It is recommended to check the fit of the abutment by radiography.
- 30. Cover the screw hole with a suitable material and cement the prosthesis with the preferred luting cement.

Warning: After the abutment screw is torqued, the abutment screw hole should be closed with suitable materials. Correct closure of the screw hole makes it possible to access the screw in cases where the abutment needs to be separated from the implant.

12. Magnetic Resonance Imaging (MRI) Safety Information

Non-clinical testing has demonstrated that Titanium Dental Materials are MR Compatible.

A patient with these system parts in his body can safely be scanned in MR systems if the following conditions are met:

Only 1.5 and 3.0 Tesla static magnetic fields

Magnetic spatial drop field up to 970 Gauss/cm (9.7 T/m) or less

Average specific absorption rate (SAR) for the whole body 2 W/kg (Normal Use Mode)

Under the above-mentioned scanning conditions, after 15 minutes of continuous scanning, our products are expected to generate a maximum temperature rise of 3°C.

13. Disposal

Disposal should be done in an environmentally sustainable manner according to local regulations. Hazardous waste and cutting/drilling tools for contaminated devices should be disposed of in suitable containers that meet special technical requirements.

14. Disclaimer of Liability

These products are part of a general concept and should only be used with the relevant original products in accordance with Mode Medikal's instructions and recommendations. The use of products not distributed by

Hazırlayan/ Prepared By
Gonca Bakırcı

Saniye Özgür

Sayfa 190 | 229



Medical Devices Regulation (EU) 2017/745 CE Technical File **Product Name: Dental Implant System Superstructures** Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 191 / 229

Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, penal or other damages arising from or in connection with any errors in professional judgment or practice related to the use of Mode Medikal products; assumes no liability, express or implied.

15.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Savfa 191 | 229



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 192 / 229

16. Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

		ngs Prepared According to th	
Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
<u> </u>	Attention	M	Production Date
(2)	Do Not Use Second Time	类	Do not expose to direct sunlight
Ţ <u>i</u>	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	Ť	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STEROLIZE	Do Not Sterilize For The Second Time

Hazırlayan/ Prepared By Gonca Bakırcı

All

Onaylayan/ Approverd By Saniye Özgür

Sog



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 193 / 229

Symbol	Symbol Description
أ	Patient name or patient ID
[31]	Date of implantation
Ų	Name and Address of the implanting healthcare institution/provider
***	Name and Address of the manufacturer
†i	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 194 / 229

Mode Abutment and Prosthetic Screws











1. Product Description

Mode abutment and prosthetic screws are part of Mode Dental Implant Systems, which is an integrated system with Mode dental implants and related abutments, cover screws, healing abutments, surgical and prosthetic parts and instruments; It is produced as a single piece using Ti6Al4V-ELI (ASTM F 136).

Mode abutment screws are implant screws that allow abutments to be fixed to implants. Mode prosthetic screws are implant screws that enable the fixation of patient-specific Geçici or permanent crowns to the relevant abutments. The abutments that Mode abutments and prosthetic screws are compatible with are shown in Table 1 below.

Table 1

Screw	Abutment	Screwdriver
	Direct Abutment	
	Esthetic Abutment	
	Esthetic Abutment 15º	
	Esthetic Abutment 25º	
Abutment Screws	Profile Abutment	
	Ti-base Non-Engaged Abutment	Mode Screwdriver
	Ti-base Engaged Abutment	
	Ti-base Cerec Abutment	
	Premill Abutment	
Prosthetic Screws	Multi Unit Abutment	
	Multi Base Abutment	

Hazırlayan/ Prepared By Gonca Bakırcı





Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 195 / 229

	NP	RP	WP
Prosthetic Screws		06.01.01.0	03
M1.4 Prosthetic Screws		06.01.01.0	04
PROVO-S Prosthetic Screws		34.00.12.0	04

	NP	RP	WP
ABUTMENT SCREW	06.0	1.01.01	06.01.01.02
M1.6- O ABUTMENT SCREW		06.01.02.	01

2. Intended Use

Mode abutment screws are intended to be used to fix Mode abutments to corresponding Mode dental implants.

Mode prosthetic screws are intended to be used to fix patient-specific Geçici or permanent crowns to the relevant abutments.

3. Target Patient Group And Intended User

Mode abutment and prosthetic screws are intended to be use with Mode dental implants and related abutments in patients with complete or partial edentulism who have completed growth and development and do not have the conditions specified in the contraindications.

Mode abutment and prosthetic screws are intended to be used by dentists only.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to use Mode dental implants safely and properly.

4. Indications

Mode abutment screws are indicated to be used to fix the abutments which are related to the prostheses to be produced for the functional, phonetic and aesthetic rehabilitation of the upper and/or lower jaw in totally or partially edentulous patients, to the placed Mode dental implants.

Mode prosthetic screws are indicated for use to fix prostheses to Mode abutments designed for the functional, phonetic and aesthetic rehabilitation of the upper and/or lower jaw in totally or partially edentulous patients.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
A	Song
· ·	



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 196 / 229

5. Contraindications

The use of Mode abutment and prosthetic screws is contraindicated in the presence of the following conditions;

- Patients medically unfit for implant treatment
- Allergy or hypersensitivity to Ti6Al4V-ELI (ASTM F 136) materials

Note: For contraindications to implant treatment, refer to the relevant Mode implant instructions for use; For the contraindications of the abutments where the screws will be used, refer to the relevant Mode abutment instructions for use.

6. Warnings / Cautions / Precautions

General

! Mode dental implant systems products should be applied by paying attention to the instructions for use provided by Mode Medikal. It is the clinician's responsibility to use the products according to these instructions and to determine whether the implants are appropriate for the individual patient situation.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

! Close collaboration between the surgeon, the prosthodontist and the dental laboratory technician is essential for successful implant treatment.

! Mode abutment and prosthetic screws should only be used in prosthesis stages with their own system elements and compatible abutments. The use of products with different brands and materials may lead to mechanical problems, failure of implants, tissue damage or aesthetic dissatisfaction.

! Mode abutment and prosthetic screws are packaged non-sterile. It is the clinician's responsibility to ensure sterility prior to use.

! Sterile handling is essential. Never use potentially contaminated components. Contamination can cause infections.

! Mode abutments and prosthetic screws are for single use and should not be reused.

! The recommended torque value is 25 Ncm for abutment screws and 15 Ncm for prosthetic screws.

! All surgical instruments and tooling used must be kept in good condition and care must be taken not to damage the implants or other components of the instruments.

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 196 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 197 / 229

Pre-operative

! A comprehensive clinical and radiological examination is necessary to determine the psychological and physical condition of the patient.

! Preoperative hard and soft tissue deficiencies may result in undesirable aesthetic results and/or negative implant angulation.

! Particular attention should be paid to patients with local or systemic factors that may affect hard tissue healing, soft tissue healing, or osseointegration of implanted implants. (eg Type 1 Diabetes, bone metabolism diseases, bleeding disorders, anticoagulant therapy, parafunctional habits such as bruxism, smoking, poor oral hygiene, pregnancy, acute periodontal diseases, orofacial radiotherapy, adjacent tissue infections, etc.)

! Particular attention should be paid to patients receiving bisphosphonate treatment.

! Implant treatment should not be applied to patients who have not completed bone growth and development.

! Placement of implants and design of prostheses should be specific to the patient. In the presence of extreme bruxism and other parafunctional habits and jaw relationship disorders, different treatment options can be planned.

During the operation

! Before placing the Mode abutment and prosthetic screws, ensure that the inner surface of the implant is clean and free from blood.

! Be careful against the risk of aspiration and swallowing due to the size of the products during intraoral application. Aspiration of products can lead to infection or unwanted physical injury.

Post-operative

! For the long-term success of treatments with the Mode dental implant system, it is important to follow up the patient regularly and to give the necessary oral hygiene training to the patient.

7. Complications And Side Effects

Dental implant treatment outcomes can be affected by many variables. Possible complications and side effects listed below may be observed after the use of Mode abutment and prosthetic screws.

- Local pain
- Micro hemorrhages
- Swelling
- Local inflammations

Onaylayan/ Approverd By Hazırlayan/ Prepared By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 198 / 229

- Gingival injuries
- Hypersensitivity and/or allergic reactions
- Other toxicity reactions
- Aspiration or swallowing of parts during operation
- Triggering of the pharyngeal reflex during the insertion of the product
- Screw fracture
- Screw loosening
- Component or prosthesis failure/fracture

8. Procedure

- I. Use the implant screw suitable for the abutment you selected for treatment. (See Table 1)
- II. Insert the Mode implant screw into the Mode screwdriver.
- III. First, tighten the screw you placed on the abutment by hand.
- IV. The Mode torque ratchet and Mode screwdriver should be used to fix the implant screws.

CAUTION: When using Mode abutment and prosthetic screws, it is recommended to ensure that the screwdriver is firmly seated on the screws to avoid aspiration/swallowing of the screws.

NOTE: The recommended torque value for tightening is 25 Ncm for abutment screws and 15 Ncm for prosthetic screws.

CAUTION: Torque of Mode abutment and prosthetic screws higher than the recommended torque values may cause screw fractures, and under-torque may cause loosening of the screws.

9. Compatibility Information

Mode abutment and prosthetic screws are compatible with Mode Medikal implant systems products and components.

Do not use implants and components made of different metals and supplied from different manufacturers with Mode dental implant system products. Mode abutments and prosthetic screws can only be used with the corresponding abutments.

10. MRI Safety Information

Mode abutment and prosthetic screws have been shown to be MRI conditional through testing in non-clinical tests. Patients using this device can be safely screened in an MRI system if the following conditions are met:

- 1.5 T and 3.0 T static magnetic field
- Maximum surface area gradient of 3,000 gauss/cm (30 T/m)

Onaylayan/ Approverd By Hazırlayan/ Prepared By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 199 / 229

 At maximum MRI system, average specific absorption rate (SAR) of the whole body 2 W/kg (Normal Operating Mode)

Under the scanning conditions described above, Mode dental implants are expected to have a maximum temperature rise of 4°C at 3.0 T and 3°C at 1.5 T after 15 minutes of continuous scanning.

In non-clinical tests, device-induced image artifact expands radially up to 2.7 cm and 2.2 cm from the implant when scanned with a gradient echo pulse array at 3.0 T and 1.5 T, respectively.

11. Cleaning And Sterilization

Mode abutment and prosthetic screws are packaged non-sterile. It is the clinician's responsibility to ensure sterility prior to use.

Sterile use is essential against the risk of contamination. Never use potentially contaminated components. Contamination can cause infections in the user and patient.

Previously used or non-sterile abutments and prosthetic screws should not be used under any circumstances.

Mode abutments and prosthetic screws can be cleaned manually or in an automatic washing unit. After cleaning, each device should be sealed and sterilized by placing it in a sterilization bag separately.

Mode Medikal recommends the following for cleaning abutment and prosthetic screws prior to use.

- I. Clean under running water by brushing inside and outside with a brush.
- II. The pre-treated product can be cleaned by hand, with ultrasonic assistance, or by an automatic cleaning method.
- III. When choosing the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

Mode Medikal recommends the following for sterilization of abutment and prosthetic screws prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121°C	Local
(444551415)	30 min.	
Pre-vacuum		

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
AK	Solog



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 200 / 229

12. Storage

The products should be stored at room temperature, in a dry place protected from direct sunlight. The products must be kept in their original packaging. Improper storage conditions may affect the product's characteristics and cause failure.

See labelings for specific storage and handling rules.

13. Disposal

Disposal of the products should be done in accordance with the rules within the framework of legal procedures and environmental requirements. Contaminated products and sharp parts are hazardous and should be disposed of as medical waste in appropriate conditions.

14. Information That Should Be Provided To The Patient

Patients should be informed about the contraindications, warnings, precautions, side effects and complications of Mode abutment and prosthetic screws.

Patients should also be informed about MR safety information.

15. Disclaimer Of Liability

These products are part of a general concept and should only be used together with the relevant original products, in accordance with Mode Medikal's instructions and recommendations. The use of products that are not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, punitive or other damages arising out of or in connection with any errors in professional judgment or application related to the use of Mode Medikal products; disclaims any liability, express or implied.

16.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Medical Devices Regulation (EU) 2017/745 CE Technical File Product Name : Dental Implant System Superstructures Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 201 / 229

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

17.Symbols

ISO 15223-1: 2021 Symbols and Meanings Prepared According to the Standard

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
(2)	Do Not Use Second Time	类	Do not expose to direct sunlight

Hazırlayan/ Prepared By
Gonca Bakırcı

Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 202 / 229

i	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
₩	Name and Address of the implanting healthcare institution/provider
~	Name and Address of the manufacturer

Hazırlayan/ Prepared By
Gonca Bakırcı

Onaylayan/ Approverd By
Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 203 / 229

***	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/Türkiye

MODE HEALING ABUTMENTS



1. Product description

Mode healing abutments are part of Mode Dental Implant Systems, which is an integrated system with Mode dental implants and related abutments, cover screws, surgical and prosthetic parts and instruments; It is produced as a single piece using Ti6Al4V-ELI (ASTM F 136) material.

Mode NP, RP, and WP healing abutments are components that support gingival healing over the implant, facilitate soft tissue shaping, and connect directly to endosseous dental implants.

Provo-S and Multi Unit-N healing abutments, on the other hand, do not connect directly to the implant but are used by establishing a connection with the implant's suprastructure.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 204 / 229

The compatibility details of Mode healing abutments narrow platform (NP), regular platform (RP), and wide platform (WP) with implants, as well as the compatibility information of Provo-S healing cap and Multi Unit-N healing cap with their respective abutments, are provided in the tables below.

Implant / Platform	Narrow Platform (NP)	Regular Platform (RP)	Wide Platform (WP)
Level İmplant			
Rapid İmplant			
Bone İmplant	NP Healing	RP Healing	WP Healing
Tissue İmplant	Abutments	Abutments	Abutments
Short implant			
Shorter İmplant			
PROVO-S IMPLANT		PROVO-S Healing Cap	

Implant / Platform	Narrow Platform (NP)	Regular Platform (RP)
Multı Unit-N Abutment	Multi Unit-N Healing Cap	

The diameter and height options of Mode narrow platform (NP), regular platform (RP), and wide platform (WP) healing abutments are provided in the table below.

Platform	NP	NP	RP	RP	RP	WP	WP	WP
Length (L)	Ø4	Ø4.5	Ø4.5	Ø5	Ø5.5	Ø5.5	Ø6	Ø6.5
2 mm	04.04.02.03	04.45.02.03	04.45.02.35	04.05.02.35	04.55.02.35	04.55.02.45	04.06.02.45	04.65.02.45
4 mm	04.04.04.03	04.45.04.03	04.45.04.35	04.05.04.35	04.55.04.35	04.55.04.45	04.06.04.45	04.65.04.45
6 mm	04.04.06.03	04.45.06.03	04.45.06.35	04.05.06.35	04.55.06.35	04.55.06.45	04.06.06.45	04.65.06.45

2. Intended Use

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 205 / 229

They are intended to used for supporting gingival healing on the implant and providing soft tissue contouring by placing them on the corresponding Mode dental implants.

3. Target Patient Group And Intended User

Mode healing abutments are intended for use with Mode dental implants in patients with complete or partial edentulism who have completed growth and development and do not have the conditions specified in the contraindications.

Mode healing abutments are intended for use by dentists only.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to use Mode dental implants safely and properly.

4. Indications

Mode healing abutments are indicated for use with Mode dental implants placed for the functional, phonetic and aesthetic rehabilitation of the upper and/or lower jaw in totally or partially edentulous patients.

5. Contraindications

The use of Mode healing abutments is contraindicated in the presence of the following conditions;

- Patients who are medically unfit for implant treatment
- Allergy or hypersensitivity to Ti6Al4V-ELI (ASTM F 136) materials

Note: For implant therapy contraindications, refer to the respective Mode implant intructions for use.

6. Warnings / Cautions / Precautions

General

! Mode dental implant systems products should be applied by paying attention to the instructions for use provided by Mode Medikal. It is the clinician's responsibility to use the products according to these instructions and to determine whether the implants are appropriate for the individual patient situation.

It is recommended that the clinician review the product-specific technique before performing the procedure. MODE MEDİCAL can provide technical information. Please contact MODE MEDICAL sales representative.

! Treatments with implants may lead to biological and mechanical problems, including bone loss and fatigue fracture of implants.

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 205 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 206 / 229

KULLANMA KILAVUZU/ USER MANUAL

! Close collaboration between the surgeon, the prosthodontist and the dental laboratory technician is essential for successful implant treatment.

! Mode healing abutments should only be used with their own system elements and paired with the correct platform during the surgical and prosthetic phases. The use of products with different brands and materials may lead to mechanical problems, failure of implants, tissue damage or aesthetic dissatisfaction.

! Mode healing abutments should be used only during the healing process. They cannot be used to support a restoration.

! Mode healing abutments are packaged non-sterile. It is the clinician's responsibility to ensure sterility prior to

! Sterile handling is essential. Never use potentially contaminated components. Contamination can cause infections.

! All surgical instruments and tooling used must be kept in good condition and care must be taken not to damage the implants or other components of the instruments.

Pre-operative

! A comprehensive clinical and radiological examination is necessary to determine the psychological and physical condition of the patient.

! Preoperative hard and soft tissue deficiencies may result in undesirable aesthetic results and/or negative implant angulation.

! Particular attention should be paid to patients with local or systemic factors that may affect hard tissue healing, soft tissue healing, or osseointegration of implanted implants. (eg Type 1 Diabetes, bone metabolism diseases, bleeding disorders, anticoagulant therapy, parafunctional habits such as bruxism, smoking, poor oral hygiene, pregnancy, acute periodontal diseases, orofacial radiotherapy, adjacent tissue infections, etc.)

! Particular attention should be paid to patients receiving bisphosphonate treatment.

! Implant treatment should not be applied to patients who have not completed bone growth and development.

! Placement of implants and design of prostheses should be specific to the patient. In the presence of extreme bruxism and other parafunctional habits and jaw relationship disorders, different treatment options can be planned.

During the operation

! Before placing the Mode healing abutments, ensure that the inner surface of the implant is clean and free from blood.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 207 / 229

! Multi Unit-N and Provo-S healing abutments are not placed directly onto the implant. They are used by connecting to the suprastructure placed on the implant.

! Be careful against the risk of aspiration and swallowing due to the size of the products during intraoral application. Aspiration of products can lead to infection or unwanted physical injury.

Post-operative

! For the long-term success of treatments with the Mode dental implant system, it is important to follow up the patient regularly and to give the necessary oral hygiene training to the patient.

7. Complications And Side Effects

Dental implant treatment outcomes can be affected by many variables. Possible complications and side effects listed below may be observed after the use of Mode healing abutments.

- Local pain
- Micro hemorrhages
- **Swelling**
- Local inflammations
- Gingival injuries
- Hypersensitivity and/or allergic reactions
- Other toxicity reactions
- Aspiration or swallowing of parts during operation
- Triggering of the pharyngeal reflex during the insertion of the product
- Growth of soft tissue on the healing abutments during the healing process
- Calculus formation on the healing abutments

8. Procedure

- I. Select the appropriate healing abutment for the implant platform and gingival height.
- II. Take the Mode healing abutment with the Mode screwdriver.
- III. Connect the healing cap to the implant you placed and tighten it by hand.
- IV. To remove the Mode healing abutment, loosen it by hand using the screwdriver.

CAUTION: When using Mode healing abutments, it is recommended to ensure that the screwdriver is firmly seated in the head to avoid the risk of aspiration/swallowing of the head.

NOTE: The recommended tightening torque is 5-10 Ncm manually by hand.

9. Compatibility Information

Mode healing abutments are compatible with Mode Medikal implant systems products and components.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 208 / 229

Do not use implants and components made of different metals and supplied from different manufacturers with Mode dental implant system products.

Mode healing abutments can only be used with those with the corresponding connection on all Mode dental implants.







03.3 - 03.7

04.1 - 04.7 - 05.2

11. MRI Safety Information

Mode healing abutments have been shown to be MRI conditional through testing in non-clinical tests. Patients using this device can be safely screened in an MRI system if the following conditions are met:

- 1.5 T and 3.0 T static magnetic field
- Maximum surface area gradient of 3,000 gauss/cm (30 T/m)
- At maximum MRI system, average specific absorption rate (SAR) of the whole body 2 W/kg (Normal Operating Mode)

Under the scanning conditions described above, Mode dental implants are expected to have a maximum temperature rise of 4°C at 3.0 T and 3°C at 1.5 T after 15 minutes of continuous scanning.

In non-clinical tests, device-induced image artifact expands radially up to 2.7 cm and 2.2 cm from the implant when scanned with a gradient echo pulse array at 3.0 T and 1.5 T, respectively.

12. Cleaning And Sterilization

Mode healing abutments are packaged non-sterile. It is the clinician's responsibility to ensure sterility prior to use.

Sterile use is essential against the risk of contamination. Never use potentially contaminated components. Contamination can cause infections in the user and patient.

Previously used or non-sterile healing abutments should not be used under any circumstances.

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 209 / 229

Mode healing abutments can be cleaned manually or in an automatic washing unit. After cleaning, each device should be sealed and sterilized by placing it in a sterilization bag separately.

Mode Medikal recommends the following for cleaning healing abutments prior to use.

- I. Clean under running water by brushing inside and outside with a brush.
- II. The pre-treated product can be cleaned by hand, with ultrasonic assistance, or by an automatic cleaning method.
- III. When choosing the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

Mode Medikal recommends the following for sterilization of healing abutments prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121°C 30 min.	Local
Pre-vacuum	55	

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

13. Storage

The products should be stored at room temperature, in a dry place protected from direct sunlight. The products must be kept in their original packaging. Improper storage conditions may affect the product's characteristics and cause failure.

See labelings for specific storage and handling rules.

14. Disposal

Disposal of the products should be done in accordance with the rules within the framework of legal procedures and environmental requirements. Contaminated products and sharp parts are hazardous and should be disposed of as medical waste in appropriate conditions.

15. Information That Should Be Provided To The Patient

Patients should be informed about the contraindications, warnings, precautions, side effects and complications of Mode healing abutments.

Patients should also be informed about MR safety information.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 210 / 229

16. Disclaimer Of Liability

These products are part of a general concept and should only be used together with the relevant original products, in accordance with Mode Medikal's instructions and recommendations. The use of products that are not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, punitive or other damages arising out of or in connection with any errors in professional judgment or application related to the use of Mode Medikal products; disclaims any liability, express or implied.

17.Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 210 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 211 / 229

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

18.Symbols

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
②	Do Not Use Second Time	类	Do not expose to direct sunlight
i	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	*	Keep away from contact with water

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 212 / 229

	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
vīv, ↓	Name and Address of the implanting healthcare institution/provider
	Name and Address of the manufacturer
Ţi —	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Shap



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 213 / 229



Explanation of unique device identifier as AIDC Format

AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/ TURKEY

MODE SINUS COVER SCREWS INSTRUCTIONS FOR USE



1. Product Description

Mode sinus cover screws are part of Mode Dental Implant Systems, an integrated system with Mode dental implants and related abutments, healing abutments, surgical and prosthetic parts and instruments; It is produced as a single piece using Ti6Al4V-ELI (ASTM F 136) material.

Mode sinus cover screws are components that cover the implant-abutment connection area during the healing process and prevent soft and hard tissue growth in this area.

While the upper part of the cover screw tightly covers the implant, the threaded part fits into the internal screw thread of the implant.

Mode sinus cover screws are designed to be compatible with all the following Mode dental implants in three different platforms, narrow platform (NP) and normal platform (RP):

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 214 / 229

Implant / Platform	Narrow Platform (NP)	Regular Platform (RP)
Level Implant		
Rapid Implant		
Bone Implant	NP Sinus Cover screw	RP Sinus Cover screw
Tissue Implant	REF: 03.02.01.03	REF: 03.02.01.35
Short Implant		
Shorter Implant		

2. Intended use

They are intended to used with the relevant Mode dental implants in the upper jaws in partially or completely edentulous patients by covering the implant-abutment connection for preventing soft and hard tissue growth in this area during the healing process area in cases where two-stage surgical technique is used and used for sinus.

3. Target Patient Group And Intended User

Mode sinus cover screws are intended for use with Mode dental implants in patients with complete or partial edentulism who have completed growth and development and do not have the conditions specified in the contraindications.

Mode sinus cover screws are intended for use by dentists only.

Clinicians must have sufficient knowledge of implantology and practice skills, as well as comply with the instructions for use, in order to use Mode dental implants safely and properly.

4. Indications

Mode sinus cover screws are indicated for use with Mode dental implants placed for the functional, phonetic and aesthetic rehabilitation of the upper jaw in totally or partially edentulous patients.

5. Contraindications

The use of Mode sinus cover screws is contraindicated in the presence of the following conditions;

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 215 / 229

- Patients who are medically unfit for implant treatment
- Allergy or hypersensitivity to Ti6Al4V-ELI (ASTM F 136) materials

Note: For implant treatment contraindications, refer to the respective Mode implant intructions for use.

6. Warnings / Cautions / Precautions

General

! Mode dental implant systems products should be applied by paying attention to the instructions for use provided by Mode Medikal. It is the clinician's responsibility to use the products according to these instructions and to determine whether the implants are appropriate for the individual patient situation.

! Treatments with implants may lead to biological and mechanical problems, including bone loss and fatigue fracture of implants.

! Close collaboration between the surgeon, the prosthodontist and the dental laboratory technician is essential for successful implant treatment.

! Mode sinus cover screws should only be used with their own system elements and paired with the correct platform during the surgical and prosthetic phases. The use of products with different brands and materials may lead to mechanical problems, failure of implants, tissue damage or aesthetic dissatisfaction.

! Mode sinus cover screws are packaged non-sterile. It is the clinician's responsibility to ensure sterility prior to use.

! Sterile handling is essential. Never use potentially contaminated components. Contamination can cause infections.

! All surgical instruments and tooling used must be kept in good condition and care must be taken not to damage the implants or other components of the instruments.

Pre-operative

! A comprehensive clinical and radiological examination is necessary to determine the psychological and physical condition of the patient.

! Preoperative hard and soft tissue deficiencies may result in undesirable aesthetic results and/or negative implant angulation.

! Particular attention should be paid to patients with local or systemic factors that may affect hard tissue healing, soft tissue healing, or osseointegration of implanted implants. (eg Type 1 Diabetes, bone metabolism diseases, bleeding disorders, anticoagulant therapy, parafunctional habits such as bruxism, smoking, poor oral hygiene, pregnancy, acute periodontal diseases, orofacial radiotherapy, adjacent tissue infections, etc.)

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 216 / 229

! Particular attention should be paid to patients receiving bisphosphonate treatment.

! Implant treatment should not be applied to patients who have not completed bone growth and development.

! Placement of implants and design of prostheses should be specific to the patient. In the presence of extreme bruxism and other parafunctional habits and jaw relationship disorders, different treatment options can be planned.

During the operation

! Before placing the Mode sinus cover screws, ensure that the inner surface of the implant is clean and free from blood.

! Be careful against the risk of aspiration and swallowing due to the size of the products during intraoral application. Aspiration of products can lead to infection or unwanted physical injury.

! After implant placement, the clinician decides whether to use immediate or delayed loading protocols, based on the evaluation of the bone quality and primary stability in the operation.

! Quality and/or quantity of remaining bone, distubed initial healing, local infections and general systemic diseases can be counted as causes of implant failures that occur immediately after the operation or after initial osseointegration has been achieved.

! Against the risk of contamination, it should be placed on the implant with the help of necessary equipment as soon as it is removed from the sterile package. Impairment of sterility may adversely affect the treatment outcome.

Post-operative

! For the long-term success of implant treatments, it is important to follow up the patient regularly and to give the necessary oral hygiene training to the patient.

7. Complications and side effects

Dental implant treatment outcomes can be affected by many variables. Possible complications and side effects listed below may be observed after the use of Mode sinus cover screws.

- Hypersensitivity and/or allergic reactions
- Other toxicity reactions
- Aspiration or swallowing of parts during operation
- Triggering of the pharyngeal reflex during the insertion of the product
- Growth of bone tissue on the sinus cover screws during the healing process
- In some cases, it becomes visible early as exposure during the healing process.

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 217 / 229

8. Procedure

- I. Take the Mode cover screw from the sterile package with the Mode screwdriver.
- II. Connect the cover screw to the implant you placed in the socket and tighten it by hand.
- III. Loosen the Mode cover screw by hand using the screwdriver to remove it.

NOTE: When using the mode sinus cover screws, it is recommended to ensure that the screwdriver is firmly seated on the screw to avoid the risk of aspiration/swallowing of the screw.

NOTE: The recommended tightening torque is 5-10 Ncm manually by hand.

9. Compatibility information

Mode sinus cover screws are compatible with Mode Medikal implant systems products and components.

Do not use implants and components made of different metals and supplied from different manufacturers with Mode dental implant system products.

Mode sinus cover screws can be used on all Mode dental implants only with the corresponding connection.



10. MRI safety information

Mode sinus cover screws have been shown to be MRI conditional through testing in non-clinical tests. Patients using this device can be safely screened in an MRI system if the following conditions are met:

- 1.5 T and 3.0 T static magnetic field
- Maximum surface area gradient of 3,000 gauss/cm (30 T/m)
- At maximum MRI system, average specific absorption rate (SAR) of the whole body 2 W/kg (Normal Operating Mode)

Under the scanning conditions described above, Mode dental implants are expected to have a maximum temperature rise of 4°C at 3.0 T and 3°C at 1.5 T after 15 minutes of continuous scanning.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Release Date:05.05.2020 Revision No: 07 Revision Date: 22.07.2024

Document No: TD.01/2.4.1

Page 218 / 229

In non-clinical tests, device-induced image artifact expands radially up to 2.7 cm and 2.2 cm from the implant

when scanned with a gradient echo pulse array at 3.0 T and 1.5 T, respectively.

21. Cleaning And Sterilization

Mode sinus cover screwss are packaged non-sterile. It is the clinician's responsibility to ensure sterility prior to use.

Sterile use is essential against the risk of contamination. Never use potentially contaminated components. Contamination can cause infections in the user and patient.

Previously used or non-sterile sinus cover screwss should not be used under any circumstances.

Mode sinus cover screwss can be cleaned manually or in an automatic washing unit. After cleaning, each device should be sealed and sterilized by placing it in a sterilization bag separately.

Mode Medikal recommends the following for cleaning sinus cover screwss prior to use.

- I. Clean under running water by brushing inside and outside with a brush.
- II. The pre-treated product can be cleaned by hand, with ultrasonic assistance, or by an automatic cleaning method.
- III. When choosing the automatic cleaning method, the appropriate detergent should be selected and the manufacturer's instructions should be followed.

Mode Medikal recommends the following for sterilization of sinus cover screwss prior to use:

Method	Conditions	Drying
Humidity temperature (autoclave)	121°C	Local
Pre-vacuum	30 min.	

Caution: Use devices immediately after sterilization. Do not store sterilized devices.

22. Storage

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 219 / 229

The products should be stored at room temperature, in a dry place protected from direct sunlight. The products must be kept in their original packaging. Improper storage conditions may affect the product's characteristics and cause failure.

See labelings for specific storage and handling rules.

23. Disposal

Disposal of the products should be done in accordance with the rules within the framework of legal procedures and environmental requirements. Contaminated products and sharp parts are hazardous and should be disposed of as medical waste in appropriate conditions.

24. Information That Should Be Provided To The Patient

Patients should be informed about the contraindications, warnings, precautions, side effects and complications of Mode sinus cover screwss.

Patients should also be informed about MR safety information.

25. Disclaimer Of Liability

These products are part of a general concept and should only be used together with the relevant original products, in accordance with Mode Medikal's instructions and recommendations. The use of products that are not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, punitive or other damages arising out of or in connection with any errors in professional judgment or application related to the use of Mode Medikal products; disclaims any liability, express or implied.

26. Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 220 / 229

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

18.Symbols

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C E 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
(2)	Do Not Use Second Time	*	Do not expose to direct sunlight
<u> </u>	Please refer to the user manual	(S)	Do not use if package is damaged

Hazırlayan/ Prepared By Onaylayan/ Approverd By Gonca Bakırcı Saniye Özgür



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 221 / 229

REF	Reference Number	Ť	Keep away from contact with water
	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
₩,	Name and Address of the implanting healthcare institution/provider
~	Name and Address of the manufacturer
!	Information website for patients

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Sog



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 222 / 229

MD	Device name
LOT	Lot Number/Batch Code
UDI	Explanation of unique device identifier as AIDC Format AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 222 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 223 / 229

MODE PROVO-S ENGAGED and NON-ENGAGED COPING INSTRUCTIONS FOR USE

1. Product description

Provo-S Engaged Coping and Provo-S Non-Engaged Coping are parts used with monoblock Provo-S dental implants and are designed to ensure the correct positioning of the implant superstructure.

Provo-S Engaged Coping: Provides a positive connection between the implant and the superstructure and increases the rotational stability of the superstructure. Especially preferred for single tooth restorations.

Provo-S Non-Engaged Coping: It is a component that allows free rotation of the superstructure within the implant. It is especially used in multiple implant restorations, bridge prostheses or multiple connections.

Provo-S engaged and non-engaged copings are manufactured using Ti6Al4V-ELI (ASTM F 136) material.

Product Name	Reference Number
Provo-S Engaged Coping	34.00.12.10
Provo-S Non-Engaged Coping	34.00.12.11

2. Intended use

These copings are used in the prosthetic stages to ensure the correct design of the implant superstructure. While Engaged Coping provides stability in single tooth restorations, Non-Engaged Coping supports passive seating of the prosthesis in multiple implant cases.

3. Target patient group and intended user

Mode dental implants are intended for use with Mode dental implants in patients with complete or partial edentulism who have completed growth and development and who do not have the conditions specified in the contraindications.

Mode dental implants may only be used by dentists.

Clinicians must have sufficient implantological knowledge and practical skills to safely and properly use Mode dental implants and must follow the instructions for use.

4. Indications

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 224 / 229

They are attached directly to the intra-osseous implants with the help of screws to provide support for implant prostheses. In completely edentulous or partially edentulous dental arches, fixed restorations provide support for prostheses.

5. Contraindications

Use is contraindicated in individuals with hypersensitivity to titanium and titanium alloys. For implant treatment contraindications, refer to the relevant Mode Medikal implant user manuals.

6. Warnings and Precautions

The patient may swallow or aspirate prosthetic parts. Care must be taken when working in the mouth. Mode medical abutments should only be used with Mode brand implants on the appropriate platform. The clinician is advised to review the product specific technique before performing the procedure. MODE MEDIKAL can provide technical information. Please contact your MODE MEDIKAL sales representative. The abutment screw should be torqued with a torque value of 25 Ncm. A torque value higher than 25 Ncm may damage the prosthetic parts. Torque values lower than the recommended value may cause abutment loosening. Prosthetic screws used in the angled Multi Unit-N Abutment series should be torqued with a torque value of 15 Ncm. Abutments are disposable devices. Mode Medikal product should be used according to the instructions for use provided by the manufacturer. It is the doctor's responsibility to use the device according to these instructions and to decide whether it is suitable for the individual patient situation. It is important that the dentist and dental technician work in harmony for a successful implant treatment. In order for implant treatment to be successful in the long term, it is important that the patient is routinely checked after the treatment is completed and oral hygiene training is provided.

7. Possible complications and side effects

The following reactions may occur during implant treatment:

- Hypersensitivity/allergic reactions
- Ingestion or aspiration of prosthetic components
- Peri-implantitis
- Failure/fracture of implant components or prosthesis
- Poor aesthetic result
- Repetition of prosthetic treatment
- Minor bleeding
- Implant loss

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 225 / 229

8. Compatible Products Information

Provo-S Engaged and Non- Engaged copings can be used with Provo-S implant diameters Ø 3.0- Ø 3.5- Ø 4.0-Ø 4.5.

Implant	İmplant Diameters
Provo- S Implant	Ø 3.0- Ø 3.5- Ø 4.0- Ø 4.5

9. Cleaning

Provo-S Engaged and Non-Engaged copings are delivered unsterilised. The products must be cleaned and sterilised before being placed in the patient's mouth.

10. Sterilization

Mode Medical copings are delivered unsterilised. Mode Medikal recommends the following for sterilisation of copings before use:

Method	Conditions	Drying
Autoclave	121°C	Local
Pre-vacuum	30 dk	

11. Procedure

a. Laboratory Procedure

- 1. Connect the impression coping with the implant analogue.
- 2. Obtain the plaster working model using soft gum silicone around the analogues.
- 3. Produce the prosthetic substructure according to the preferred method. The substructure can be produced by casting, CAD/CAM or metal sintering.
- **4.** Produce the final prosthesis according to the preferred laboratory technique.

Hazırlayan/ Prepared By	Onaylayan/ Approverd By
Gonca Bakırcı	Saniye Özgür
At	Song



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 226 / 229

27. Disposal

Disposal of the products should be done in accordance with the rules within the framework of legal procedures and environmental requirements. Contaminated products and sharp parts are hazardous and should be disposed of as medical waste in appropriate conditions.

28. Information That Should Be Provided To The Patient

Patients should be informed about the contraindications, warnings, precautions, side effects and complications of Mode sinus cover screwss.

Patients should also be informed about MR safety information.

29. Disclaimer Of Liability

These products are part of a general concept and should only be used together with the relevant original products, in accordance with Mode Medikal's instructions and recommendations. The use of products that are not distributed by Mode Medikal and made by third parties will void any expressed or implied warranty or other obligations of Mode Medikal.

Mode Medikal products should be used in accordance with the instructions for use provided by the manufacturer. It is the clinician's responsibility to use products in accordance with these instructions and to determine whether the product is appropriate for the individual patient situation. The clinician is also responsible for regularly following the latest updates on the relevant Mode Medikal product and its applications. Mode Medikal is not responsible for any direct, indirect, punitive or other damages arising out of or in connection with any errors in professional judgment or application related to the use of Mode Medikal products; disclaims any liability, express or implied.

30. Additional Information

Non Sterile products do not have a specified shelf life.

Life Time

The useful life of the products is determined as 10 years.

The SSCP link of the products is in the link below;

https://modeimplant.com/images/SSCP-Dental-Implant-Systems.pdf

NOTICE REGARDING SERIOUS INCIDENT

Hazırlayan/ Prepared By Gonca Bakırcı

Onaylayan/ Approverd By Saniye Özgür

Sayfa 226 | 229



Ürün Adı: Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 227 / 229

For a patient/user/third party in the European Union and in countries with an identical regulatory regime (Regulation 2017/745/EU on Medical Devices); if, during the use of this device or as a result of its use, a serious incident has occurred, please report it to the manufacturer and to your national authority. The contact information for the manufacturer of this device to report a serious incident is as follows:

Manufacturer Name: Mode Medikal Sanayi ve Ticaret Limited Şirketi

Adress: Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa/İstanbul

Phone: 0212 612 64 09

Mail: info@modeimplant.com

Related Person: Melis AKYÜZ

Mail: melis@modemed.com

Phone: 0553 373 36 42

18.Symbols

Symbol	Explanation of the Symbol	Symbol	Explanation of the Symbol
C € 2696	Notified Body Number	NON	Non Steril
\triangle	Attention	M	Production Date
(2)	Do Not Use Second Time	类	Do not expose to direct sunlight
[]i	Please refer to the user manual		Do not use if package is damaged
REF	Reference Number	Ť	Keep away from contact with water

Hazırlayan/ Prepared By Gonca Bakırcı



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020 Revision No: 07

Revision Date: 22.07.2024

Page 228 / 229

	Barcode Number	MR	A product that has been proven safe in the MR environment under defined conditions.
MD	Medical Device	LOT	Lot Number
121°C	Sterilisation temperature	STERRIZE	Do Not Sterilize For The Second Time

Symbol	Symbol Description
† ?	Patient name or patient ID
[31]	Date of implantation
vīv, ↓	Name and Address of the implanting healthcare institution/provider
	Name and Address of the manufacturer
P 1	Information website for patients
MD	Device name
LOT	Lot Number/Batch Code

Hazırlayan/ Prepared By Gonca Bakırcı Onaylayan/ Approverd By Saniye Özgür

Shop



Ürün Adı : Dental İmplant Sistemi Üstyapıları

KULLANMA KILAVUZU/ USER MANUAL

Document No: TD.01/2.4.1 Release Date:05.05.2020

Revision No: 07 Revision Date: 22.07.2024

Page 229 / 229



Explanation of unique device identifier as AIDC Format

AIDC: Automatic identification and data capture format (e.g. linear or 2D-Barcodes)

Yenidoğan Mah. Abdi İpekçi Cad. No:58 Bayrampaşa,İstanbul/TURKEY

Hazırlayan/ Prepared By Gonca Bakırcı

All